

Describing a Quantum Channel by State Tomography of a Single Probe State

CHANG-SHUI YU ^(a), HE-SHAN SONG

School of Physics and Optoelectronic Technology, Dalian University of Technology, Dalian 116024, P. R. China

PACS 03.65.Ud – Entanglement and quantum nonlocality
 PACS 42.50.Dv – Quantum state engineering and measurements
 PACS 03.65.Ta – Foundations of quantum mechanics; measurement theory

Abstract. - A general law is presented for (composite) quantum systems which directly describes the time evolution of quantum states (with one or both components) through an arbitrary noisy quantum channel. It is shown that the time evolution of all quantum states through a quantum channel can be completely captured by the evolution of a single 'probe state'. Thus in order to grasp the information of the final output states subject to a quantum channel, especially an unknown one, it only requires quantum state tomography of a single probe state, which dramatically simplifies the practical operations in experiment.

Quantum states are the basic carrier of quantum information [1]. The core of all the quantum information processing (QIP) including quantum communication [2] and quantum computation [3] is the controlled time evolution of quantum state in essence [4]. However, in realistic scenario, quantum states will be unavoidably and greatly disturbed by the undesired coupling to the uncontrolled degree of freedom usually termed as 'environment' and described as a 'quantum channel'. As a consequence, besides the state itself the valuable properties of quantum states such as coherence [5,6], entanglement [7,8] of composite systems and so on will be greatly corrupted. The precise characterization of some properties of quantum states usually largely relies on the evaluation of quantum states, if these properties such as entanglement does not correspond to a direct observable for a general unknown quantum state [9-11]. Furthermore, quantum channel is not restricted to the previous interaction between the system and environment. It is a general notion of any a input/output device governed by quantum mechanics including the controlled interactions, for example, the dynamical action of a quantum gate in a quantum computer etc [12]. Therefore, it is of practical importance to precisely explore the time evolution of quantum states on which a reliable QIP task depends.

In general cases, there is no direct way to evaluate the time evolution of quantum states. One has to begin with

considering the dynamics of 'system of interests + environment' governed by quantum principle [13-16]. It is implied that the concrete description of the quantum channel has been known by the quantum process tomography that includes a series of quantum state tomography [17] and is usually quite complex [3]. In experiment, the time evolution of quantum states could be described by determining the initial and final states in terms of quantum state tomography no matter whether the quantum channel is known. However, it is a drawback that the procedure needs to be repeated every time with different input states chosen. In the present Letter, we provide a direct and general scheme in terms of the evolution of a given probe state to describe the evolution of quantum states of an arbitrary (composite) quantum system which (the components of which) undergoes an arbitrary (especially unknown) quantum channel. The distinguished advantages of our scheme are as follows: 1) Especially for unknown quantum channels, it is only necessary to do state tomography of the single probe state, stead of repeating the same procedure for different input states or doing quantum process tomography (In other words, it is not necessary to know the concrete description of quantum channel). 2) The scheme can be directly applied to any quantum mechanical input/output process. Thus all information of the final states can be learned and the properties of interests such as coherence or entanglement etc. can be obtained by a sequent simple calculation [18,19].

Let us first consider an $(N \otimes N)$ -dimensional bipartite

^(a)quaninformation@sina.com; ycs@dlut.edu.cn

quantum states ρ_0 which can be expanded in a representation spanned by maximally entangled states given by

$$|\Phi_j\rangle = \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} e^{i\frac{2j_0 k \pi}{N}} |k\rangle |k \oplus j_1\rangle, j = Nj_0 + j_1, \quad (1)$$

where $j_0, j_1 = 0, 1, \dots, N-1$, $|k\rangle$ is the computational basis and ' $\oplus$ ' denotes the addition modulo N . Suppose each subsystem of ρ_0 undergoes a quantum channel represented by $\$1$ and $\$2$, respectively, then the final state can be given by $\rho_f = (\$1 \otimes \$2) \rho_0 / p$, where $p = \text{Tr}[(\$1 \otimes \$2) \rho_0]$ is the joint probability for channels $\$1$ and $\$2$ which corresponds to non-trace-preserving channels [20]. In the representation of maximally entangled states, ρ_f can be expanded as

$$\rho_f = \frac{1}{p} \sum_{mn} (\$1 \otimes \$2) [|\Phi_m\rangle \langle \Phi_m| (\mathbf{1} \otimes \$2) \rho_0 |\Phi_n\rangle \langle \Phi_n|], \quad (2)$$

with $\mathbf{1}$ being the identity. Throughout the letter, we refer to $(N \times N)$ matrix ψ (i.e. without $|\rangle\rangle$) as the matrix notation of any pure state $|\psi\rangle = \sum_{i,j=0}^{N-1} a_{ij} |ij\rangle$ with matrix elements $\langle i|\psi|j\rangle = a_{ij}$. For a maximally entangled state $|\Phi_m\rangle$, it hence follows that [21]

$$|\Phi_m\rangle = (\Phi_m P^{-1} \otimes \mathbf{1}) |P\rangle, \quad (3)$$

where $|P\rangle = \sum_{i,j=0}^{N-1} \tilde{a}_{ij} |ij\rangle$, called 'probe quantum state', is a generic entangled pure state with full-rank P (which can be explicitly written as

$$P = \begin{pmatrix} \tilde{a}_{00} & \tilde{a}_{01} & \cdots & \tilde{a}_{0(N-1)} \\ \tilde{a}_{10} & \tilde{a}_{11} & \cdots & \tilde{a}_{1(N-1)} \\ \vdots & \vdots & \ddots & \vdots \\ \tilde{a}_{(N-1)0} & \tilde{a}_{(N-1)1} & \cdots & \tilde{a}_{(N-1)(N-1)} \end{pmatrix},$$

or can be directly obtained by the method provided above (3).) and P^{-1} denotes the inverse matrix of P . Φ_m in (3) are simple unitary transformations determined by (1). For example, for the state of a pair of qubits, Φ_m corresponds to the three Pauli matrices and the identity, respectively. Thus $\rho_0 = \sum_k p_k (\phi_k P^{-1} \otimes \mathbf{1}) |P\rangle \langle P|$

$\times ([P^\dagger]^{-1} \phi_k^\dagger \otimes \mathbf{1})$ for a potential decomposition $\{p_k, \phi_k\}$. Based on Jamiołkowski isomorphism [22], (2) can be rewritten as

$$\begin{aligned} \rho_f &= \frac{1}{p} \sum_{mnk} p_k \langle \Phi_m | (\phi_k P^{-1} \otimes \mathbf{1}) [(\mathbf{1} \otimes \$2) |P\rangle \langle P|] \\ &\quad \times ([P^\dagger]^{-1} \phi_k^\dagger \otimes \mathbf{1}) | \Phi_n \rangle (\$1 \otimes \mathbf{1}) [|\Phi_m\rangle \langle \Phi_n|] \\ &= \frac{p_{II} p_{III}}{p} \sum_{mn} \text{Tr} \left\{ S (P^{-1} \otimes \Phi_m^*) \left[\frac{(\mathbf{1} \otimes \$2) |P\rangle \langle P|}{p_{II}} \right] \right. \\ &\quad \times ([P^\dagger]^{-1} \otimes \Phi_n^T) S \rho_0^* \left. \right\} (\mathbf{1} \otimes \Phi_m^T [P^T]^{-1}) \\ &\quad \times \left[\frac{(\$1 \otimes \mathbf{1}) |P\rangle \langle P|}{p_I} \right] (\mathbf{1} \otimes [P^*]^{-1} \Phi_n^*), \end{aligned} \quad (4)$$

where S is the swapping operator defined as $S|j\rangle|k\rangle = |k\rangle|j\rangle$, ' T ' denotes transpose operation, $p_I = \text{Tr}[(\$1 \otimes \mathbf{1}) |P\rangle \langle P|]$, $p_{II} = \text{Tr}[(\mathbf{1} \otimes \$2) |P\rangle \langle P|]$ and p is the normalization factor of (4). Once we select one probe state $|P\rangle$, the evolution of ρ_0 will completely be determined by the two output probe states $\frac{(\$1 \otimes \mathbf{1}) |P\rangle \langle P|}{p_I}$

and $\frac{(\mathbf{1} \otimes \$2) |P\rangle \langle P|}{p_{II}}$ up to several fixed operations which are independent of the initial input states and quantum channels. In experiment, so long as one does state tomography of the final states of the probe state corresponding to the two quantum channels, one can directly obtain the final output state of ρ_0 . Thus (4) provides a general, direct and simple relationship between the input state ρ_0 and the output state ρ_f . Quantum channel's action on the initial quantum state has been completely captured by a probe quantum state $|P\rangle$ which is alternative depending on the experimental convenience. The maximally entangled

state $|\Phi\rangle = \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} |k\rangle |k\rangle$ is a special probe state with which (4) has a simple form by replacing P by $\mathbf{1}$. Note that it is not necessary to employ the same probe state $|P\rangle$ for both channels $\$1$ and $\$2$.

The direct relationship (4) can be reduced to describe the time evolution of an N -dimensional quantum state ϱ_0 through a quantum channel $\$$. To derive the explicit expression of the time evolution, we first introduce an auxiliary N -dimensional quantum state σ which is completely a mathematical skill to double the original quantum state space. Thus the bipartite joint quantum state can be written as $\rho'_0 = \varrho_0 \otimes \sigma$ and the corresponding final state $\rho'_f = \frac{1}{p_s} \$\varrho_0 \otimes \sigma$ with $p_s = \text{Tr}(\$ \varrho_0)$, which is equivalent to the case of one-sided quantum channel in (4) and can be directly obtained by setting $\rho_0 = \rho'_0$, $\$1 = \$$ and $\$2 = \mathbf{1}$. Hence the output final state corresponding to ϱ_0 is given by $\varrho_f = \text{Tr}_a \rho'_f$ with Tr_a denoting trace over the auxiliary system. It is obvious that the final state ϱ_f is independent of the auxiliary state σ . Repeating the same procedure to (4), one can obtain a compact form of ϱ_f :

$$\begin{aligned} \varrho_f &= \frac{p_{III}}{p_s} \sum_{mn} \langle \Phi_m | \frac{(\$ \otimes \mathbf{1}) |P\rangle \langle P|}{p_{III}} | \Phi_n \rangle \\ &\quad \times \Phi_m P^{-1} \varrho_0 (P^{-1})^\dagger \Phi_n^\dagger, \end{aligned} \quad (5)$$

where $p_{III} = \text{Tr}((\mathbb{I} \otimes \mathbb{I}) |P\rangle \langle P|)$. (5) does not include the auxiliary state itself as expected, hence σ can be arbitrary quantum states and does not work in practical experiment. The output state given in (5) is completely determined by the one-sided quantum channel's actions on the $(N \otimes N)$ -dimensional probe quantum states. (5) provides a general input/output relationship for a given quantum state through an arbitrary quantum channel.

In fact, (5) is a general law suitable for all quantum states no matter whether the quantum systems under consideration are composite (including multipartite quantum states) or not. However it is of the most importance that the auxiliary Hilbert space introduced should be completely consistent with that under consideration, unless the quantum channel is a nonlocal one acting on all components of the composite system which can equivalently be mapped to a single quantum system. For example, if we consider an $(N_1 \otimes N_2)$ -dimensional bipartite quantum state through a one-sided or two-sided quantum channel, the auxiliary should also be an $(N_1 \otimes N_2)$ -dimensional bipartite quantum state. Thus, the probe quantum state should be an $[(N_1 N_2) \otimes (N_1 N_2)]$ -dimensional generic bipartite entangled state which is a quadripartite quantum state in essence. However, if both components of the composite system undergo a common nonlocal quantum channel, only a genuine bipartite probe state will be enough.

Let us finally compare (4) and (5) briefly when we consider an $(N \otimes N)$ -dimensional bipartite quantum state. In (4), the probe states are two separate generic $(N \otimes N)$ -dimensional entangled states. However, in (5) one has to prepare a generic $(N^2 \otimes N^2)$ -dimensional entangled state (a quadripartite entangled state) as the probe state which is obviously not as easy to implement in experiment [23] as that in (4). However, we again would like to emphasize that (5) is a universal law for all quantum states.

In conclusion, we have presented a direct and general input/output law for the time evolution of quantum states with the system or its components through an arbitrary quantum channel. It has shown that quantum channel's action on any quantum states can be completely captured by a preconditioned probe quantum state. In other words, for any quantum channel, especially for an unknown one, it is enough to only explore the time evolution of one of the probe states instead of repeating the same procedure for every potential initial input state. Thus the output probe state, just like a quantum gate, directly relates the input and output states by which all information on the final output states can be learned.

This work was supported by the National Natural Science Foundation of China, under Grant No. 10805007 and No. 10875020, and the Doctoral Startup Foundation of Liaoning Province.

REFERENCES

- [1] C. H. Bennett and D. P. DiVincenzo. *Nature* **404**, 247 (2000).
- [2] D. Bouwmeester, J.-W. Pan, K. Mattle, M. Eibl, H. Weinfurter and A. Zeilinger. *Nature* **390**, 575 (2000).
- [3] M. A. Nielsen, and I. L. Chuang. *Quantum Computation and Quantum Information* (Cambridge Univ, Press, Cambridge, 2000).
- [4] M. Steffen, M. Ansmann, B. C. Bialczak, N. Katz, E. Lucero, R. McDermott, M. Neeley, E. M. Weig, A. N. Cleland and J. M. Martinis. *Science* **313**, 1423-1425 (2006).
- [5] W. H. Zurek. *Phys Today* **44**, 36 (1991).
- [6] M. Schlosshauer. *Rev. Mod. Phys.* **76**, 1267 (2004).
- [7] J. H. Eberly and T. Yu. *Science* **316**, 555 (2007).
- [8] M. P. Almeida, F. Melo, M. Hor-Meyll, A. Salles, S. P. Walborn, P. H. S. Ribeiro and L. Davidovich. *Science* **316**, 579 (2007).
- [9] P. Horodecki. *Phys. Rev. Lett.* **90**, 167901 (2003).
- [10] S. P. Walborn, P. H. S. Ribeiro, L. Davidovich, F. Mintert, and A. Buchleitner. *Nature* **440**, 1022 (2006).
- [11] Chang-shui Yu and He-shan Song. *Phys. Rev. A* **76**, 022324 (2007).
- [12] C. Roos. *Nature physics* **4**, 97 (2008).
- [13] C. W. Gardiner and P. Zoller. *Quantum noise* (Springer-Verlag, Berlin, 1991).
- [14] M. B. Plenio, P. L. Knight. *Rev. Mod. Phys.* **70**, 101 (1998).
- [15] M. O. Scully and M. S. Zubairy. *Quantum optics* (Cambridge Univ, Press, Cambridge, 1997).
- [16] T. Yu and J. H. Eberly. *Phys. Rev. Lett.* **97**, 140403 (2006).
- [17] H. Häffner, et al. *Nature* **438**, 643 (2005).
- [18] T. Yu and J. H. Eberly. *Phys. Rev. B* **68**, 165322 (2003).
- [19] W. K. Wootters. *Phys. Rev. Lett.* **80**, 2245 (1998).
- [20] T. Konrad, F. Melo, M. Tiersch, C. Kasztelan, A. Aragao and A. Buchleitner. *Nature physics* **4**, 99 (2008).
- [21] P. Horodecki and R. Horodecki. *Quantum Information and Computation* **1**, 45 (2001).
- [22] A. Jamiołkowski. *Rep. Math. Phys.* **3**, 275 (1972).
- [23] C. -Y. Lu, X.-Q. Zhou, O. Gühne, W. B. Gao, J. Zhang, Z. -S. Yuan, A. Goebel, T. Yang and J. -W. Pan. *Nature Physics* **3**, 91 (2007).